

Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 0 505 047 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention
of the grant of the patent:
24.04.1996 Bulletin 1996/17

(51) Int Cl.⁶: **C07D 305/14, C07D 317/70,
A61K 31/335**

(21) Application number: **92301652.1**

(22) Date of filing: **27.02.1992**

(54) **Rearranged taxol compounds and method of using in testing of in vivo activity**

Umlagerungsverbindungen von Taxol und deren Verwendung bei der Bestimmung von in vivo-Aktivitäten

Dérivés de Taxol redispesés et procédé d'utilisation dans l'essai d'activité in vivo

(84) Designated Contracting States:
**AT BE CH DE DK ES FR GB GR IT LI LU MC NL
PT SE**

(30) Priority: **18.03.1991 US 670432**

(43) Date of publication of application:
23.09.1992 Bulletin 1992/39

(73) Proprietor: **VIRGINIA TECH INTELLECTUAL
PROPERTIES, INC.
Blacksburg, VA 24060 (US)**

(72) Inventors:
• **Kingston, David G.I.
Blacksburg, VA 24060 (US)**
• **Samaranyake, Gamini
Blacksburg, VA 24060 (US)**

(74) Representative: **Pattullo, Norman et al
Murgitroyd and Company
373 Scotland Street
Glasgow G5 8QA (GB)**

(56) References cited:
• **JOURNAL OF THE AMERICAN CHEMICAL
SOCIETY**, vol. 93, no. 9, 5 May 1971,
WASHINGTON, DC, US, pages 2325 - 2327; M.C.
WANI ET AL.: 'Plant antitumour agents. VI. The
isolation and structure of taxol, a novel
antileukaemic and antitumour agent from
Taxus brevifolia'
• **JOURNAL OF ORGANIC CHEMISTRY**, vol. 56,
no. 17, 16 August 1991, WASHINGTON, DC, US,
pages 5114 - 5119; G. SAMARANAYAKE ET
AL.: 'Modified taxols. 5. Reaction of taxol with
electrophilic reagents and preparation of
rearranged taxol derivative with tubulin
assembly activity'
• **The Chemistry of Taxol, Pharmac. Ther. Vol. 52,**
pp. 1-34, 1991

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 0 505 047 B1

Description

FIELD OF THE INVENTION

This application is directed to rearranged taxol compounds, and more particularly is directed to taxol derivatives with an opened oxetane ring, taxol derivatives in which the oxetane ring is opened and the A-ring is contracted, and taxol derivatives with a contracted A-ring.

BACKGROUND OF THE INVENTION

The naturally occurring diterpenoid taxol has great potential as an anti-cancer drug, and has shown activity in several tumor systems. Background information on taxol, the mechanisms of taxol's biological activity, and the synthesis of water soluble derivatives from taxol, are described in co-pending application Serial Number 07/573,731, filed August 28, 1990. All references cited herein are incorporated by reference as if reproduced in full below.

Taxol is in short supply and is relatively expensive. Further, total synthesis of taxol is quite difficult due to the bridged bicyclic portion of the taxane ring structure. Therefore, it is highly desirable to find taxol derivatives with similar biological activities, which lend themselves to easier total synthesis than taxol.

There is also a need for a method to quickly determine the biological activities of new compounds or pharmaceutical compounds having bioactivities or structures similar to taxol. The short supply and expense of taxol makes impractical the use of taxol as a standard in determining the bioactivities of other compounds; thus, it is highly desirable that a range of other standards with known biological activities be available to determine the bioactivity of taxol derivatives and other compounds relative to taxol. Useful standards should be derivatives of taxol, or the standards should be compounds which have similar structures to taxol, but which are more readily available or which can be synthesized easier than taxol. At present, some derivatives, which do not exhibit the same high biological activity as taxol, are thrown away; this waste would be eliminated by a method which uses taxol derivatives, which have significantly less biological activity than taxol, as standards in bioactivity testing, rather than utilizing more taxol, which is already in short supply and very expensive.

Thus, there is a need for taxol derivatives having a range of *in vivo* and *in vitro* activities, and there is a need for taxol derivatives or compounds having similar biological activities to taxol which are capable of easier total synthesis than taxol.

OBJECTS OF THE INVENTION

Thus, it is a primary object of the present invention to synthesize taxol derivatives having varying *in vitro* and *in vivo* activities for use as standards in determining the bioactivity of other compounds in comparison to taxol.

It is a further object of the present invention to synthesize derivatives of taxol, having similar *in vivo* activities, which are capable of easier total synthesis than taxol.

SUMMARY OF THE INVENTION

These and other objects of the present invention are accomplished through synthesis of compounds useful for the aforementioned purposes and objectives. In a preferred embodiment, taxol is treated with mesyl chloride to prepare a taxol derivative with a contracted A-ring, which has comparable activity to taxol in a tubulin depolymerization assay, and which shows cytotoxicity against KB cells in a cell culture assay. In an alternate preferred embodiment, taxol is treated with triethyloxonium tetrafluoroborate (Meerwein's reagent) to produce a taxol derivative with an opened oxetane ring. In another alternate preferred embodiment, reaction of taxol with acetyl chloride yields a taxol derivative in which the oxetane ring is opened and the A-ring is contracted. All of the aforementioned products show *in vivo* activity in KB cell culture assays. Further, the preferred compounds have different *in vivo* activities, which makes them ideal to form a range of standards for biological testing of other compounds.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a generalized representation of a taxol compound structure.

Figure 2 is a generalized representation of a taxol compound with an opened oxetane ring.

Figure 3 is a generalized representation of a taxol compound having a contracted A-ring and an opened oxetane ring.

Figure 4 is a generalized representation of a taxol compound with a contracted A-ring.

Figure 5 is a representation of structure of an acetonide of taxol.

Figure 6 is an alternate generalized representation of a taxol compound.

Figure 7 is a generalized representation of the hydrogenated form of the compound illustrated in Figure 3.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

With reference to Figure 1, the structure of taxol (1) is represented when R1 is C(O)CH₃, R2 is OH, R3 is H, and R4 is H. The derivatives of taxol discussed herein are represented in Figures 1-7, with specific structure variables, R1-R7, defined in Table I below. For the sake of convenience, structures will be referred to by the structure numbers provided in Table I.

TABLE I

Variable Definitions For Structures Discussed Herein And Illustrated In Figures 1-7		
Structure Number	Figure Number	Variable Definitions
(1)	1	R1=C(O)CH ₃ , R2=OH, R3=H, R4=H
(2)	1	R1=C(O)CH ₃ , R2=H, R3=OH, R4=H
(3)	1	R1=H, R2=H, R3=OH, R4=H
(4)	2	R1=C(O)CH ₃ , R2=OH, R3=H, R4=H, R5=H, R6=H, R7=C(O)CH ₃
(5)	2	R1=C(O)CH ₃ , R2=OC(O)CH ₃ , R3=H, R4=C(O)CH ₃ , R5=H, R6=H, R7=C(O)CH ₃
(6)	2	R1=C(O)CH ₃ , R2=OC(O)CH ₃ , R3=H, R4=C(O)CH ₃ , R5=C(O)CH ₃ , R6=H, R7=C(O)CH ₃
(7)	3	R1=C(O)CH ₃ , R2=OC(O)CH ₃ , R3=H, R4=C(O)CH ₃ , R5=OC(O)CH ₃ , R6=OH, R7=OC(O)CH ₃
(8)	7	R1=C(O)CH ₃ , R2=OC(O)CH ₃ , R3=H, R4=C(O)CH ₃ , R5=OC(O)CH ₃ , R6=OH, R7=OC(O)CH ₃
(9)	4	R1=C(O)CH ₃ , R2=OSi(CH ₂ CH ₃) ₃ , R3=Si(CH ₂ CH ₃) ₃
(10)	4	R1=C(O)CH ₃ , R2=OH, R3=H
(11)	6	R1=C(O)CH ₃ , R2=OSi(CH ₂ CH ₃) ₃ , R3=OH, R4=Si(CH ₂ CH ₃) ₃
(12)	5	R1=C(O)CH ₃ , R2=C(O)CH ₃
(13)	6	R1=C(O)CH ₃ , R2=OSi(CH ₂ CH ₃) ₃ , R3=OSO ₂ CH ₃ , R4=Si(CH ₂ CH ₃) ₃

Due to the complexity of the taxol structure, and the incomplete understanding of the relationship between the taxol structure and its activity, it was difficult or impossible to predict how derivatives of taxol would behave; further, the complexity of the taxol structure made prediction of reaction product structures equally difficult. The present invention overcomes these problems through the synthesis of taxol derivatives with a contracted A-ring (Figure 4), with a contracted A-ring and an opened oxetane ring (Figure 3), or with an opened oxetane ring (Figure 2), which retain *in vivo* activity, although at different levels than taxol.

The taxol derivative having a contracted A-ring structure, otherwise known in a preferred embodiment as A-Nortaxol (10), is easier to perform total synthesis of than taxol since the bridged bi-cyclic portion of the diterpenoid structure has been fused. Further, it has been discovered that A-Nortaxol has an *in vitro* activity very close to that of taxol in a tubulin depolymerization assay. Thus, A-Nortaxol (10) can be used in a similar fashion to taxol (1).

It is believed that the oxetane ring (the D-ring in taxol (1)) is susceptible to ring-opening by reaction with electrophilic reagents, so initial experiments were performed using zinc bromide as an electrophile to attempt to open the oxetane ring on taxol. Taxol (1) was combined with zinc bromide in methanol at ambient temperature, but the oxetane ring was not opened, and the taxol simply underwent epimerization of the C-7 hydroxyl group and cleavage of the 10-acetyl group to yield 7-epitaxol (2) and 10-deacetyl-7-epitaxol (3).

The relatively mild conditions created by zinc bromide in methanol were selected because it was believed that the more vigorous conditions created by other electrophiles and other solvents may result in taxol derivatives without any substantial biological activity due to destruction of part or all of the bioactive portions of the molecule. However, it was surprisingly discovered that treatment of taxol with different electrophilic groups resulted in rearrangement of different portions of the taxol structure, and that some of the various products retained *in vivo* activity. More specifically, reaction of taxol with the strong electrophile triethyloxonium tetrafluoroborate (Meerwein's reagent) resulted in a taxol derivative (4) with an opened oxetane ring. Reaction of taxol with acetyl chloride yielded a product (7) in which the oxetane ring was opened, but the A-ring was contracted. Treatment of the protected taxol (11) with mesyl chloride, followed by deprotection, resulted in a taxol derivative with a contracted A-ring alone, A-Nortaxol (10).

It was also surprisingly discovered that these rearranged taxol derivatives (4), (7), and (10) all retain varying degrees

of in vivo activity, as determined by KB cell culture assays. A-Nortaxol (10) also showed very similar activity to taxol in a tubulin depolymerization assay, and showed higher in vivo activity than the acetyl chloride product (7) and the Meerwein product (4). Thus, the range of activity shown by the taxol derivatives of the present invention make them ideal for use as standards in biological testing. Further, A-nortaxol (10) and the acetyl chloride product (7) are easier to perform synthesis on than taxol since the A and B rings are fused in the products rather than bridged (note that the bridge bond between A and B rings in taxol is extremely difficult to synthesize). Thus, the present invention includes a method, which is highly useful to researchers in determining the activity of compounds in comparison to taxol (1) and taxol derivatives; the method has an additional advantage, since it is not necessary to use the limited and expensive supply of taxol as a control or standard.

In a preferred process, new compounds or production line pharmaceutical compounds, which must be tested to ensure bioactivity in comparison to taxol, are subjected to bioactivity evaluation in parallel with standards utilizing compounds (4), (7), and (10). Since the relative bioactivities of products (4), (7), and (10) are known (in comparison to one another as well as to taxol), the relative bioactivity of the compound being tested to taxol can be determined in a reliable fashion. Thus, in addition to the in vivo activity of compounds (4), (7), and (10), which makes the compounds useful as cytotoxic agents, they are also useful in extending the limited supply of taxol. This is because the method of the present invention replaces taxol with new compounds in testing other compounds which are used as cytotoxic agents in general, or, which are used as anti-cancer or anti-leukemic drugs in humans.

METHODS AND MATERIALS

Specific reaction methods are described in more detail in the following non-limiting examples. The methods used herein are generally described in the Journal Of Organic Chemistry, 51, pp. 797-802 (1986). Low resolution mass spectrometry data were obtained on a VG 7070 E-HF mass spectrometer. Exact mass measurements were performed at the Midwest Center for Mass Spectrometry, NSF Regional Instrumentation Facility (Grant Che-8211164). The term "standard work-up" in the following non-limiting examples includes extraction with a suitable solvent (usually ethyl acetate or methylene chloride), washing the extract with water, drying over magnesium sulfate or sodium sulfate, and evaporation in vacuo. All technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art. Other methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention.

EXAMPLE 1

Taxol (1) was reacted with zinc bromide, $ZnBr_2$ as follows: 100 mg taxol (0.117 mmol) was combined with 3.3 g $ZnBr_2$ (0.41 mmol) in 10 ml $CHCl_3$:MeOH(1:4) and stirred for 24 hours at 40° C. Water was added to the mixture, and the standard work-up (as defined above in "Methods") yielded a white solid. The solid was purified by chromatography to obtain two compounds, which were identified as 7-epi-taxol (2) and 10-deacetyl-7-epitaxol (3) by comparison to published data (see Journal of Natural Products, 49, pp. 665-669 (1986), and Journal of Natural Products, 44, pp. 312-319 (1981); compounds (2) and (3) have been previously described by McLaughlin et al. in Journal of Natural Products, 44, pp. 312-319 (1981). Thus, it was necessary to try more vigorous reaction conditions to open the oxetane ring, despite the risk that the more vigorous conditions would cause other portions of the taxol structure to be altered to an extent that the resulting compound would lack any substantial bioactivity.

EXAMPLE 2

Taxol (1) was reacted with Meerwein's reagent as follows: A solution of taxol in dichloromethane was prepared by adding 100 ml taxol (0.117 mmol) to dry dichloromethane. The taxol solution in dichloromethane was cooled and stirred while 200 μ l triethyloxonium tetrafluoroborate (1 M in CH_2Cl_2) was added drop-wise from a freshly opened bottle. These conditions were maintained for 30 minutes, at which time the reaction was quenched with 3 ml of ethereal HCl (1:2 mixture of 1N HCl:ether) followed by stirring for 10 minutes. Standard work-up gave a crude solid, which was further purified by flash chromatography and PTLC to yield 53 mg (51%) of a taxol compound with an opened oxetane ring (4) (The "Meerwein product"). NMR, MS, and IR (KBr) were performed on a sample of the Meerwein product (4) and its melting point was determined, with the characterization data and NMR data presented in Tables II and III below:

TABLE II

Characterization Data For Product (4) (Resulting From Reaction Of Taxol With Meerwein's Reagent)	
Melting Point	160-164°C (amorphous solid)
IR (KBr) cm^{-1}	1745(s), 1670(m), 1535(w), 1505(w), 1474(w), 1395(m), 1120(m), 1080(m), 1060(m)
MS(FAB) m/z	872 (MH^+ , 100), 854 (MH^+ , H_2O)
High-resolution mass spectrum	calculated (MH^+) 872.3493; observed 872.3463

TABLE III

**NMR Data For Reaction Product Of
Taxol With Meerwein's Reagent, $\text{C}_{47}\text{H}_{54}\text{NO}_{15}$ (4)**

Carbon Position	^1H Shift (PPM from TMS) Coupling Hertz
2	5.56 (d, 6)
3	4.03 (3, 6)
5	3.70 (br s)
6	a
7	4.49 (dd 4, 11)
10	6.57 (s)
13	6.01 (br dd, 4, 11)
14	2.45 (dd, 11, 16)
	3.08 (dd, 4, 16)
15	
16	1.12 (s)
17	1.12 (s)
18	2.10 (s) ^b
19	1.22 (s)
20	3.85 (ABq, 11 $\Delta\nu_{AB} = 86$)
2'	4.70 (br s)
3'	5.92 (dd, 2, 9)
NH	7.19 (br d, 9)
OAc	1.65 (s)
	2.25 (s)
2-OBz	8.03 (m)
	7.3-7.6 (m) ^c
3'-NBz	8.03 (m)
	7.3-7.6 (m) ^c
3'-Ph	7.3-7.6 (m) ^c
OH	3.91 (s)

a peak concealed under signals from methyl group

b determined by decoupling experiment

c overlapping peaks

EXAMPLE 3

The Meerwein product (4) was acetylated as follows: 5 mg of the Meerwein product (4) (0.0057 mmol) was dissolved

in 100 μ l of pyridine, and 300 μ l of acetyl chloride (excess) was added to the solution; the solution was kept at room temperature for 30 minutes, and warmed to 60° C for one hour, followed by quenching with water. Standard work-up yielded crude material which was purified on PTLC to yield 3 mg (55%) of chromatographically homogeneous 2',7-diacetyl-D-seco-taxol (5). Structure was confirmed with ¹H NMR.

EXAMPLE 4

2',7-diacetyl-D-seco-taxol (5) was acetylated as follows: 8 mg 2',7-diacetyl-D-seco-taxol (5) (0.008 mmol) was dissolved in 750 μ l tetrahydrofuran (THF) and to this solution dicyclohexylcarbodiimide (5 mg, 2.5 eq) 4 μ l acetic anhydride (5 eq) and a catalytic amount of pyrrolidino pyridine were added. The stirred solution was heated to 60° C for 7.5 hours, the solvent was then evaporated and the residue extracted into ethyl acetate. Standard work-up yielded a crude mixture which was purified by PTLC with 1% MeOH/CHCl₃ to yield 3 mg (38% yield at 63% conversion) of 2',5,7-triacetyl-D-seco-taxol (6). Characterization data is presented in Table IV and NMR data is presented in Table V below.

TABLE IV

Characterization Data For 2', 5, 7-Triacetyl-D-Seco-Taxol (6)	
IR cm ⁻¹	1740(s), 1720(s), 1676(m), 1625(s), 1225(s)
MS(FAB)m/z (relative intensity)	998(MH ⁺ , 13) 980(MH ⁺ , 12) 936(6) 848(23), 650(100).

TABLE V

NMR Data For 2', 5, 7-Triacetyl-D-Seco-Taxol (6)

Carbon Position	¹ H Shift (PPM from TMS) Coupling Hertz
2	5.63 (d, 5)
3	4.04 (d, 5)
5	5.26 (m) ^c
6	a
7	5.43 (br d, 9)
10	6.42 (s)
13	6.03 (m)
14	2.40 (dd, 9, 15)
	3.07 (dd, 5, 15)
15	
16	1.13 (s)
17	1.11 (s)
18	2.24 (s)
19	1.38 (s)
20	4.01 (ABq, 12 $\Delta\nu_{AB}=57$)
2'	5.26 (m) ^c
3'	6.12 (dd, 3, 10)
NH	7.20 (br d, 9)
OAc	1.98 (s), 2.13 (s), 2.16 (s), 2.19 (s), 2.20 (s)
2-OBz	8.20 (m)
3'-NBz	7.8 (m)

3'-Ph
OH7.2-7.6 (m)^c

- 5 **a** peak concealed under signals fr m methyl group
 b determined by decoupling experiment
 c overlapping peaks

EXAMPLE 5

10

The Meerwein product (4) was then treated with 2,2-dimethoxypropane as follows: 6 mg of the Meerwein product (4) (0.007 mmol) and 200 μ l 2,2-dimethoxypropane was combined in 500 μ l dry dichloromethane along with a catalytic amount of p-toluenesulfonic acid, and stirred for one hour. Standard work-up yielded a crude product, which was further purified by PTLC to obtain 6 mg (95%) of pure acetonide (12) (Figure 5). The structure of the acetonide (12) was confirmed by ¹H NMR and by mass spectrometry; MS(FAB) m/z (relative intensity): 916 (MNa⁺, 100), 855 (MNa⁺-HOAc-H, 25), 832 (MNa⁺-CH₃CO-C₃H₅, 50), 761(40). ¹H NMR data is presented in Table VI below.

15

TABLE VI

20

NMR Data For Acetonide (12)
Of The Meerwein Product (4)

25

Position ¹H Shift (PPM from TMS)
 Coupling Hertz

30

2	5.34 (d, 8)
3	3.06 (d, 8)
5	4.34 (m)

35

40

45

50

55

6	6 _{ax} 1.95 (ddd 5, 11, 15) ^b 6 _{eq} 2.36 (ddd 2, 5, 15)
7	4.48 (dd, 5, 11)
10	6.38 (d, 2)
13	5.68 (d, 2)
14	2.57 (AB part of ABX, 9, 14 $\Delta\nu_{AB}=62$)
15	
16	4.67 (s), 4.75 (s)
17	1.63 (s)
18	1.67 (s) ^b
19	1.62 (s)
20	4.15 (ABq, 12, $\Delta\nu_{AB}=26$)
2'	4.50 (d, 3)
3'	5.60 (dd, 3, 8)
NH	6.98 (br d, 8)
OAc	1.83 (s), 2.17 (s)
2-OBz	8.10 (m)
25	7.3-7.4 (m) ^c
3'-NBz	7.73 (m)
	7.3-7.4 (m) ^c
3'-Ph	7.3-7.4 (m) ^c
OH	
30	Other 1.30 (s) ^d 1.33 (s)

a peak concealed under signals from methyl group

b determined by decoupling experiment

c overlapping peaks

d signals of the methyl groups of the acetone

EXAMPLE 6

Taxol (1) was reacted with acetyl chloride under the following conditions: 200 mg taxol (0.23 mmol) was dissolved in 2 ml acetyl chloride, and the solution was refluxed for one hour. The reaction was quenched with ice water and ethyl acetate, and stirred for 30 minutes. Standard work-up yielded a white solid. Recrystallization of the white solid from ethyl acetate and hexanes yielded acetylated A-Nor-D-seco-taxol (7) (See Figure 3) as white needles (156 mg, 68%). Characterization data for acetylated A-Nor-D-seco-taxol (7) is provide in Table VII and NMR data is provided in Table VIII below.

TABLE VII

Characterization Data For Acetylated A-Nor-D-seco-taxol (7)	
Melting Point	140-142°C
IR(CHCl ₃) cm ⁻¹	1750(s), 1660(m), 1606(m), 1372(m), 1282(m), 1156(m)
MS(FAB)m/z	1002(MNa ⁺ , 35), 676(MNa ⁺ -side-chain, 15), 616 (675-HOAc, 30), 554 (676-PhCOOH, 20), 494(616-PhCOOH, 30), 411(24), 372(4), 177(100)
High resolution mass spectrum	calculated for C ₅₃ H ₅₇ NO ₁₇ Na(MNa ⁺)=1002.3524; observed 1002.3557

TABLE VIII

5 **NMR Data For Acetylated A-Nor-D-seco -tax 1 (7)**

Carbon Position	¹ H Shift (PPM from TMS) Coupling Hertz
10 2	5.35 (d,7)
3	3.54 (d,7)
15 5	5.28 (br s)
6	a
7	5.54 (dd,4,13)
10	6.38 (s)
13	5.67 (t,7)
20 14	2.64 (m)
15	
16	4.69 (s), 4.82 (s)
17	1.62 (s)
18	1.82 (s)
25 19	1.53 (s)
20	4.15 (ABq, 12 Δv _{AB} =55)
2'	5.52 (d,2)
3'	6.02 (dd,2,9)
30 NH	7.03 (d,9)
OAc	1.85 (s), 2.00 (s), 2.13 (s), 2.17 (s), 2.22 (s)
35 2-OBz	7.93 (m), 7.2-7.9 (m) ^c
3'-NBz	7.85 (m) 7.2-7.9 (m) ^c
3'-Ph	7.2-7.9 (m) ^c
40 OH	3.75 (s)
Other	

a peak concealed under signals from methyl group

b determined by decoupling experiment

c overlapping peaks

45 **EXAMPLE 7**

50 The Meerwein product (4) was treated with acetyl chloride as follows: 50 mg of the Meerwein product (4) (0.057 mmol) was dissolved in CHCl₃; 3 mg pyrrolidinopyridine (an amount sufficient to catalyze the reaction) and excess acetyl chloride (5eq) were added to the solution. Excess triethylamine was then added dropwise to the stirred solution at room temperature. A red color developed, which disappeared when more acetyl chloride was added; additional triethylamine and acetyl chloride were added until a single major product was obtained. The reaction was stopped after a total reaction time of 5 hours by addition of 3 ml of water and stirring for 30 minutes. Standard work-up was followed with an additional

55 wash with 3N HCl. The mixture of products was subjected to PTLC to obtain the major component as 10 mg of a white solid (18%). ¹H NMR, mass spectrometry, and infrared spectrometry confirmed that this compound was identical to the acetyl product of Example 6, acetylated A-Nor-D-seco-taxol (7)

EXAMPLE 8

The acetylated A-Nor-D-seco-taxol (7) was hydrogenated to its dihydro derivative (8) (Figure 7) as follows: 24 mg of the acetyl chloride product (7) (0.023 mmol) was dissolved in 2.5 ml ethyl acetate, and hydrogenated over Pd/H₂. After 24 hours, the catalyst was filtered off, and the solvent was evaporated to yield a crude solid, which consisted of product and unreacted starting material; the unreacted starting material was not separable from the product. The crude product was dissolved in methylene chloride and treated with 5 mg m-chloroperbenzoic acid (58%) at room temperature for 3 hours; this converted the starting material to its separable epoxide. The solvent was evaporated and the residue subjected to PTLC with 4% MeOH/CHCl₃ to yield 8 mg pure hydrogenated product (8) (35%) along with 11 mg of a mixture of diastereomeric epoxides. The hydrogenated product (8) was recrystallized from ethyl acetate and hexanes. Characterization data is presented in Table I below and NMR data is presented in Table IX below.

TABLE IX

Characterization Data for Hydrogenated Acetylated A-Nor-D-Seco-Taxol (8)	
Melting Point	148-150° C
IR (KBr) cm ⁻¹	1740(s), 1720(m), 1640(m), 1220(m), 910(m)
MS(FAB) m/z (relative intensity)	1004(MNa ⁺ , 100), 962(MNa ⁺ -C ₃ H ₆ , 10), 944(MNa ⁺ -HOAc, 15)

TABLE X

5 **NMR Data for Hydr generated
Acetylated A-Nor-D-Seco-Taxol (8)**

Carbon Position	¹ H Shift (PPM from TMS) Coupling Hertz
2	5.17 (d, 8)
3	3.68 (d, 8)
5	4.94 (br d, 11)
6	a
7	5.54 (dd, 5, 11)
10	6.36 (s)
13	5.72 (m)
14	2.48 (AB part of ABX, 14, 16 $\Delta\nu_{AB}=65$)
15	1.60 (m)
16	0.76 (d, 7)
17	0.78 (d, 7)
18	1.83 (s)
19	1.53 (s)
20	4.11 (ABq, 11 $\Delta\nu_{AB}=78$)
2'	5.40 (d, 3)
3'	5.95 (dd, 3, 8)
NH	7.02 (d, 8)
OAc	1.17 (s), 1.18 (s), 2.00 (s), 2.13 (s), 2.17 (s)
2-OBz	7.91 (m), 7.2-7.6 (m) ^c
3'-NBz	7.83 (m) 7.2-7.6 (m) ^c
3'-Ph	7.2-7.6 (m) ^c
OH	
Other	

a peak concealed under signals from methyl group
 b determined by decoupling experiment
 c overlapping peaks

EXAMPLE 9

50 Taxol was reacted with imidazole and triethylsilylchloride as follows: 238 mg of solid imidazole (10 eq) was added to a solution of 200 mg taxol (0.234 mmol) in 2.5 ml DMF; 196 μ l triethylsilylchloride (10 eq) was then added to the stirred solution while the solution was at room temperature, followed by warming the solution to 45-50°C. After 2 hours, the solution was diluted with water, and extracted with ethyl acetate. The crude solid obtained after evaporation of the solvent was purified on a silica gel flash column to yield 242 mg (96%) of pure 2',7-bis (triethylsilyl)taxol (11) (Figure 6). Characterization data is presented in Table XI and NMR data is presented in the Table XII below.

TABLE XI

Characterization Data For 2',7-bis(triethylsilyl)taxol (11)	
Melting Point	122-123°C
IR cm ⁻¹	1740(s), 1720(s), 1660(s), 1640(m), 1240(s), 810(m)
MS(FAB) m/z (relative intensity)	1104(MNa ⁺ , 100), 1003(30), 981(MNa ⁺ -PhCOOH, 10)

TABLE XII

NMR Data For 2',7-bis(triethylsilyl)taxol (11)

Carbon Position	¹ H Shift (PPM from TMS) Coupling Hertz
2	5.71 (m)
3	3.88 (d, 6)
5	4.94 (br d, 11)
6	a
7	4.48 (dd, 6, 14)
10	6.44 (s)
13	6.20 (m)
14	a
15	
16	1.22 (s)
17	1.18 (s)
18	2.02 (s)
19	1.70 (s)
20	4.27 (ABq, 11 Δν _{AB} =63)
2'	4.71 (d, 3)
3'	5.71 (m)
NH	7.12 (d, 9)
OAc	2.17 (s), 2.55 (s)
2-OBz	8.12 (m)
3'-NBz	7.2-7.5 (m) ^c
3'-Ph	7.2-7.5 (m) ^c
OH	7.2-7.5 (m) ^c
Other	0.48 (6H, m) ^c 0.57 (6H, m) 0.81 (9H, t, 8) 0.91 (9H, t, 8)

a peak concealed under signals from methyl group
b determined by decoupling experiment
c overlapping peaks

EXAMPLE 10

2',7-bis(triethylsilyl)taxol (11) was treated with triethylamine and mesyl chloride (methane sulfonyl chloride) as fol-

lows: 30 mg of 2',7-bis(triethylsilyl)taxol (11) (0.028 mmol) was added to 3 ml of dry methylene chloride, and cooled to -15°C under argon. 600 µl of triethylamine (154 eq) was added, followed by addition of 300 µl of mesyl chloride (138 eq) in 1 ml methylene chloride, with the additions made over a 5 minute period. The mixture was allowed to warm to -5 to 0° C, and this temperature was maintained for a total reaction time of 2.5 hours, at which time 50% conversion of the starting material to 2',7-bis(triethylsilyl)-A-Nortaxol (9) (Figure 4) was observed. The solution was cooled again to -15° C, and 1 ml triethylamine and 500 µl mesyl chloride were added; this latter procedure was repeated one additional time. The reaction was then stopped by adding 2 ml triethylamine, 5 ml water, and 5 ml ethyl acetate. Standard work-up yielded crude material, which was purified by PTLC to give 6 mg (20%) of 2',7-bis(triethylsilyl)-A-Nor-taxol (9) (Figure 4) along with 2 mg starting material and 2 mg 7-triethylsilyl-A-Nor-taxol. It is believed that the mesylate (13) (Figure 6) is formed as an intermediate between (11) and (9). Mass spectrometry data, m/z, are as follows: 1086(MNa⁺, 45), 1064 (MH⁺, 75), 1005(MH⁺-OAc, 25), 975(MH⁺-OAc-CH₂O, 15), 963(MH⁺-OAc-C₃H₈, 15), 820(MH⁺-PhCOOH, PhCONH₂-H, 100). NMR data is presented in Table XIII below.

TABLE XIII

15

NMR Data For 2',7-bis(triethylsilyl)-A-Nor-taxol(9)

Carbon Position	¹ H Shift (PPM from TMS) Coupling Hertz
2	5.54 (d, 8)
3	3.53 (d, 8)
5	5.02 (d, 8)
6	1.90 (m)
7	4.53 (dd, 8, 5)
10	6.39 (s)
13	5.81 (br t, 7)
14	2.40 (m)
	2.60 (m)
15	
16	4.66 (s), 4.75 (s)
17	1.65 (s)
18	1.62 (s)
19	1.73 (s)
20	4.15 (ABq, 12) Δν _{AB} =26)
2'	4.60 (d, 2)
3'	5.64 (dd, 2, 11)
NH	7.15 (d, 11)
OAc	2.02 (s), 2.40 (s)
2-OBz	8.01 (m)
	7.2-7.5 (m) ^c
3'-NBz	7.70 (m)
	7.2-7.5 (m) ^c
3'-Ph	7.2-7.5 (m) ^c
OH	
Other	0.38 (6H, m) ^a 0.60 (6H, m) 0.75 (9H, t, 8) 0.90 (9H, t, 8)

55

- a peak concealed under signals from methyl group
 b determined by decoupling experiment
 c overlapping peaks
 e signals of the ethyl groups of the TES groups

EXAMPLE 11

2',7-bis(triethylsilyl)-A-Nor-taxol (9) was treated with pyridinium hydrofluoride as follows: 67 ml 2',7-bis(triethylsilyl)-A-Nor-taxol (9) (0.07 mmol) was dissolved in 1 ml dry THF under argon. The solution was cooled to 0°C and 100 µl pyridinium hydrofluoride (70% in pyridine) was added. After 3 hours the cooling bath was removed, and the reaction was allowed to proceed for an additional 45 hours at room temperature. The reaction was quenched with 2 ml aqueous pyridine (10% v/v pyridine). Standard work-up resulted in a crude solid which was purified by PTLC with 8% Me-OH/CHCl₃, as the eluent. A-nor-taxol (10) was obtained as a white solid in 55% yield (29 mg). (Note that deprotection of the silyl ether (9) yielded a mixture of products when tetrabutylammonium fluoride was used instead of pyridinium hydrofluoride.) Mass spectrometry data, m/z (relative intensity), as follows: 836(MH⁺, 100), 776(MH⁺-HOAc, 30), 551 (836-side-chain-H, 10), 307(20); High resolution mass spectrum calculated for C₄₇H₅₀NO₁₃(MH⁺) was 836.3282, observed 836.3272. ¹H NMR is presented in Table XIV below.

TABLE XIV**¹H NMR Data For A-Nor-Taxol (10)**

Carbon Position	¹ H Shift (PPM from TMS) Coupling Hertz
2	5.49 (d, 8)
3	3.48 (d, 8)
5	5.04 (d, 8)
6	1.86 (dd, 11, 15); 2.59 (ddd, 8, 9, 15)
7	4.63 (dd, 9, 11)
10	6.32 (s)
13	5.71 (m) ^c
14	2.04 (dd, 8, 13) 2.42 (dd, 13, 6)
15	
16	4.69 (br s) 4.76 (br s)
17	1.59 (s)
18	1.61 (s)
19	1.64 (s)
20	4.24 (ABq, 8, Δν _{AB} =34)
2'	4.67 (d, 2)
3'	5.71 (dd, 11, 2) ^c
NH	6.89 (d, 11)
OAc	2.17 (s), 2.36 (s)
2-OBz	8.10 (m) 7.2-7.6 (m) ^c
3'-NBz	7.66 (m) 7.2-7.6 (m) ^c
3'-Ph	7.2-7.6 (m) ^c
OH	
Other	
a peak concealed under signals from methyl group	
b determined by decoupling experiment	
c overlapping peaks	

BIOLOGICAL TESTING OF TAXOL DERIVATIVES

Taxol (1), the Meerwein product (4), the acetyl chloride product (7), and A-nor-taxol (10) were tested in KB cell culture assays. All of the samples showed in vivo activity as shown in Table XV below (note that an ED₅₀ (μg/ml) of less than 4.0 indicates in vivo activity). Taxol has been shown to have great potential as an anti-cancer drug, so that these compounds can be used in a similar fashion to taxol as cytotoxic agents.

TABLE XV

Bioactivity Of Modified Taxols In KB Cell Culture	
Compound	ED ₅₀ (μg/ml) cell culture
Taxol (1)	0.00001
Meerwein product (4)	2.3
Acetyl Chloride product (7)	2.5
A-Nortaxol (10)	2.0

Taxol (1), the Meerwein product (4), the acetyl chloride product (7), and A-Nor-Taxol (10) were also subjected to tubulin depolymerization assays with the results presented in Table XVI below.

TABLE XVI

In Vitro Activity In Tubulin Depolymerization Assay	
Compound	ID ₅₀ (μM) in tubulin depolymerization assay
Taxol (1)	0.3
Meerwein product (4)	>6.3
A-Nortaxol (10)	0.9

Note that A-Nor-Taxol (10) has an activity which is very similar to taxol in the tubulin depolymerization assay. Since the A-Nor-Taxol (10) structure has a fused A-ring, A-Nor-Taxol (10) is a more likely candidate for total synthesis than taxol. Further, because A-Nor-taxol has been found to be cytotoxic, and to have an in vitro activity close to taxol's in a tubulin depolymerization assay, it is believed that A-Nor-taxol (10) may have bioactivity mechanisms very similar to taxol's (1).

With reference to Table XV, note that the in vivo activity of A-Nor-Taxol (10), the Meerwein product (4), and the acetyl chloride product (7) are 2.0, 2.3, and 2.5 respectively. This range of in vivo activities makes these compounds useful as standards for testing other compounds, eliminating the need for taxol standards. Since, the relative in vivo activity of these compounds is now known, the bioactivity of an unknown compound can be determined with respect to these compounds without having to refer to any published data or without use of taxol as a standard.


A preferred process for testing the bioactivity of unknown compounds, or for testing taxol-derived pharmaceuticals, includes treatment of an unknown sample in a bioactivity test, and simultaneous treatment under the same conditions of at least one, preferably of all three, of the following products: A-Nor-Taxol (10), the Meerwein product (4), and the acetyl chloride product (7). By comparison of results for the unknown sample with the results for the products (4), (7), and (10), the in vivo activity of the unknown can be determined.

Due to the bioactivity demonstrated for some of the compounds of the present invention, it is believed that compounds (4)-(12) will be useful directly as, or as intermediates for, antineoplastic, antileukemic and anticancer prodrugs or drugs, and their use as such is considered to be a part of the present invention.

Contemplated equivalents of the rearranged taxol compounds of the present invention include rearranged taxol compounds having a contracted A-ring, an opened oxetane ring, or a contracted A-ring and an opened oxetane ring, which have one or more side chain or ring substituents substituted with a group such as, but not limited to, substitution of -H, -OH, -OR, -NR, -Ar, -OC(O)CH₃, Ph-O-, Ph-N-, or =O for another non-interfering moiety.


Claims

1. A rearranged taxol compound having the following structure




wherein R1 is $\text{C}(\text{O})\text{CH}_3$, R2 is H or $\text{Si}(\text{CH}_2\text{CH}_3)_3$, and R3 is H or $\text{Si}(\text{CH}_2\text{CH}_3)_3$.

20 2. A taxol compound having the following structure:



wherein R1 is $\text{C}(\text{O})\text{CH}_3$, R2 is $\text{OSi}(\text{CH}_2\text{CH}_3)_3$, R3 is OSO_2CH_3 , and R4 is H or $\text{Si}(\text{CH}_2\text{CH}_3)_3$.

40 3. A rearranged taxol compound having the following structure:




wherein R1 is H or $\text{C}(\text{O})\text{CH}_3$, R2 is H or $\text{C}(\text{O})\text{CH}_3$.

4. A process for producing taxol compounds with an A-ring contracted to five members, comprising the steps of:
 - (a) protecting hydroxylated positions on taxol; and
 - (b) reacting said protected taxol with a mesyl halide.
5. A process according to Claim 4, further comprising the step of deprotecting said protected taxol to form the compound claimed in Claim 1 wherein R2 is H and R3 is H.
6. A process for producing taxol compounds with an A-ring contracted to five members, comprising the steps of:
 - (a) reacting taxol with a trialkylsilyl halide to form a taxol silyl ether; and
 - (b) reacting taxol silyl ether with a mesyl halide.
7. A process according to claim 6, further comprising the step of: (c) following step (b), removing silyl groups on said taxol silyl ether to yield the compound claimed in claim 1 wherein R2 is H and R3 is H.
8. The process of Claim 7, wherein step (c) is carried out by reacting said silylated compound claimed in Claim 1 wherein R2 is H and R3 is H, with pyridinium hydrofluoride.
9. The process of Claim 6, 7 or 8, wherein said steps (a) and (b) are carried out in basic conditions.
10. The process of any one of Claims 6 to 9, wherein said trialkylsilyl halide is triethylsilyl chloride, and said mesyl halide is mesyl chloride.
11. A process for producing the compound of Claim 3, comprising the reaction of taxol with triethyl oxonium tetrafluoroborate, and reacting the resulting compound with dimethoxypropane in the presence of p-toluenesulfonic acid.
12. A cytotoxic agent comprising a taxol compound having an A-ring contracted to five members.
13. The cytotoxic agent of Claim 12, wherein said taxol compound has an intact oxetane D-ring.


Patentansprüche

1. Taxol-Umlagerungsverbindung mit der folgenden Struktur:




worin R1 = C(O)CH₃, R2 = H oder Si(CH₂CH₃)₃ und R3 = H oder Si(CH₂CH₃)₃.

2. Taxolverbindung mit der folgenden Struktur:



worin $R1 = C(O)CH_3$, $R2 = OSi(CH_2CH_3)_3$, $R3 = OSO_2CH_3$ und $R4 = H$ oder $Si(CH_2CH_3)_3$.

- 20 3. Taxol-Umlagerungsverbindung mit der folgenden Struktur:



worin $R1 = H$ oder $C(O)CH_3$, $R2 = H$ oder $C(O)CH_3$.

- 40 4. Verfahren zur Herstellung von Taxolverbindungen mit einem A-Ring, der zu fünf Gliedern kontrahiert ist, bestehend aus den Stufen:

- (a) Schützen der hydroxylierten Positionen an Taxol; und
- 45 (b) Reagieren des geschützten Taxols mit einem Mesylhalid.

5. Verfahren nach Anspruch 4, welches die weitere Stufe eines Befreiens des geschützten Taxols zur Ausbildung der in Anspruch 1 beanspruchten Verbindung umfaßt, worin $R2 = H$ und $R3 = H$.

- 50 6. Verfahren zur Herstellung von Taxolverbindungen mit einem A-Ring, der zu fünf Gliedern kontrahiert ist, bestehend aus den Stufen:

- (a) Reagieren von Taxol mit einem Trialkylsilylhalid zur Ausbildung eines Taxolsilylethers; und
- 55 (b) Reagieren des Taxolsilylethers mit einem Mesylhalid.


7. Verfahren nach Anspruch 6, welches weiter die Stufe (c) im Anschluß an die Stufe (b) umfaßt, die aus einem Entfernen der Silylgruppen an dem Taxolsilylether besteht, um die in Anspruch 1 beanspruchte Verbindung zu erhalten,

worin R2 = H und R3 = H.

8. Verfahren nach Anspruch 7, worin die Stufe (c) ausgeführt wird durch ein Reagieren der in Anspruch 1 beanspruchten silylierten Verbindung, worin R2 = H und R3 = H, mit Pyridiniumhydrofluorid.
9. Verfahren nach Anspruch 6, 7 oder 8, worin die Stufen (a) und (b) unter basischen Bedingungen durchgeführt werden.
10. Verfahren nach einem der Ansprüche 6 bis 9, worin das Trialkylsilylhalid Triethylsilylchlorid ist und das Mesylhalid Mesylchlorid ist.
11. Verfahren zur Herstellung der Verbindung nach Anspruch 3, bestehend aus der Reaktion von Taxol mit Triethyloxoniumtetrafluorborat und einem Reagieren der resultierenden Verbindung mit Dimethoxypropan in der Gegenwart von p-Toluolsulfonsäure.
12. Cytotoxisches Mittel, bestehend aus einer Taxolverbindung mit einem A-Ring, der zu fünf Gliedern kontrahiert ist.
13. Cytotoxisches Mittel nach Anspruch 12, worin die Taxolverbindung einen intakten Oxetan D-Ring hat.


Revendications

1. Dérivé de taxol redispósé ayant la structure suivante :




dans laquelle R1 est C(O)CH₃, R2 est H ou Si(CH₂CH₃)₃, et R3 est H ou Si(CH₂CH₃)₃.

2. Dérivé de taxol ayant la structure suivante :



dans laquelle R1 est $C(O)CH_3$, R2 est $OSi(CH_2CH_3)_3$ et R3 est OSO_2CH_3 , et R4 est H ou $Si(CH_2CH_3)_3$.

20 3. Dérivé de taxol redispesé ayant la structure suivante :



dans laquelle R1 est H ou $C(O)CH_3$, R2 est H ou $C(O)CH_3$.

40 4. Procédé pour produire des dérivés de taxol avec un anneau A contracté à cinq membres, comprenant les étapes de :

- (a) protéger les positions hydroxylées sur le taxol ; et
- (b) faire réagir ledit taxol protégé avec un halogénure de mésyle.

45 5. Procédé selon la revendication 4, comprenant en outre l'étape de déprotéger ledit taxol protégé pour former le dérivé revendiqué dans la revendication 1 dans lequel R2 est H et R3 est H.

6. Procédé de production de dérivés de taxol avec un anneau A contracté à cinq membres, comprenant les étapes de :

- (a) faire réagir le taxol avec un halogénure de trialkylesilol pour former un éther silol taxol ; et
- (b) faire réagir l'éther silol taxol avec un halogénure de mésyle.

7. Procédé selon la revendication 6, comprenant en outre l'étape de : (c) après l'étape (b), enlever les groupes silol sur ledit éther silol taxol pour obtenir le dérivé revendiqué dans la revendication 1 dans lequel R2 est H et R3 est H.

55 8. Procédé selon la revendication 7, dans lequel l'étape (c) est réalisée en faisant réagir ledit composé silolé revendiqué dans la revendication 1 dans lequel R2 est H et R3 est H, avec de l'hydrofluorure de pyridine.

EP 0 505 047 B1

9. Procédé selon les revendications 6, 7 ou 8, dans lequel lesdites étapes (a) et (b) sont réalisées en milieu basique.
10. Procédé selon l'une quelconque des revendications 6 à 9, dans lequel ledit halogénure de trialkylesilol est le chlorure de trialkylesilol, et ledit halogénure de mésyle est le chlorure de mésyle.
- 5 11. Procédé de production d'un dérivé selon la revendication 3, comprenant la réaction du taxol avec le tétrafluoroborate de triéthyle oxonium, et la réaction du dérivé qui en résulte avec le diméthoxypropane en présence d'acide p-toluènesulfonique.
- 10 12. Agent cytotoxique comprenant un dérivé de taxol ayant un anneau A contracté à cinq membres.
13. Agent cytotoxique de la revendication 12, dans lequel ledit dérivé de taxol a un anneau D oxétane intact.

15

20

25

30

35

40

45

50

55




FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) Publication number:

0 505 047 A1

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 92301652.1

(51) Int. Cl.⁵: C07D 305/14, C07D 317/70,
A61K 31/335

(22) Date of filing: 27.02.92

(30) Priority: 18.03.91 US 670432

(43) Date of publication of application:
23.09.92 Bulletin 92/39

(84) Designated Contracting States:
AT BE CH DE DK ES FR GB GR IT LI LU MC
NL PT SE

(71) Applicant: VIRGINIA TECH INTELLECTUAL
PROPERTIES, INC.
1900 Kraft Drive, Suite 107
Blacksburg, VA 24060(US)

(72) Inventor: Kingston, David G.I.
1409 Locust Avenue
Blacksburg, VA 24060(US)
Inventor: Samaranyake, Gamini
1303 Glade Road
Blacksburg, VA 24060(US)

(74) Representative: Pattullo, Norman et al
Murgitroyd and Company Mitchell House 333
Bath Street
Glasgow G2 4ER(GB)

(54) Rearranged taxol compounds and method of using in testing of in vivo activity.

(57) In a preferred embodiment, taxol is treated with mesyl chloride to prepare a taxol derivative with a contracted A-ring, which has comparable activity to taxol in a tubulin depolymerization assay, and which shows cytotoxicity against KB cells in a cell culture assay. In an alternate preferred embodiment, taxol is treated with triethyloxonium tetrafluoroborate (Meerwein's reagent) to produce a taxol derivative with an opened oxetane ring. In another alternate preferred embodiment, reaction of taxol with acetyl chloride yields a taxol derivative in which the oxetane ring is opened and the A-ring is contracted. All of the aforementioned products show in vivo activity in KB cell culture assays. Further, the preferred compounds have different in vivo activities, which makes them ideal to form a range of standards for use in biological testing of other compounds.




FIG. 1

FIELD OF THE INVENTION

This application is directed to rearranged taxol compounds, and more particularly is directed to taxol derivatives with an opened oxetane ring, taxol derivatives in which the oxetane ring is opened and the A-ring is contracted, and taxol derivatives with a contracted A-ring.

BACKGROUND OF THE INVENTION

The naturally occurring diterpenoid taxol has great potential as an anti-cancer drug, and has shown activity in several tumor systems. Background information on taxol, the mechanisms of taxol's biological activity, and the synthesis of water soluble derivatives from taxol, are described in co-pending application Serial Number 07/573,731, filed August 28, 1990. All references cited herein are incorporated by reference as if reproduced in full below.

Taxol is in short supply and is relatively expensive. Further, total synthesis of taxol is quite difficult due to the bridged bicyclic portion of the taxane ring structure. Therefore, it is highly desirable to find taxol derivatives with similar biological activities, which lend themselves to easier total synthesis than taxol.

There is also a need for a method to quickly determine the biological activities of new compounds or pharmaceutical compounds having bioactivities or structures similar to taxol. The short supply and expense of taxol makes impractical the use of taxol as a standard in determining the bioactivities of other compounds; thus, it is highly desirable that a range of other standards with known biological activities be available to determine the bioactivity of taxol derivatives and other compounds relative to taxol. Useful standards should be derivatives of taxol, or the standards should be compounds which have similar structures to taxol, but which are more readily available or which can be synthesized easier than taxol. At present, some derivatives, which do not exhibit the same high biological activity as taxol, are thrown away; this waste would be eliminated by a method which uses taxol derivatives, which have significantly less biological activity than taxol, as standards in bioactivity testing, rather than utilizing more taxol, which is already in short supply and very expensive.

Thus, there is a need for taxol derivatives having a range of in vivo and in vitro activities, and there is a need for taxol derivatives or compounds having similar biological activities to taxol which are capable of easier total synthesis than taxol.

OBJECTS OF THE INVENTION

Thus, it is a primary object of the present invention to synthesize taxol derivatives having varying in vitro and in vivo activities for use as standards in determining the bioactivity of other compounds in comparison to taxol.

It is a further object of the present invention to synthesize derivatives of taxol, having similar in vivo activities, which are capable of easier total synthesis than taxol.

SUMMARY OF THE INVENTION

These and other objects of the present invention are accomplished through synthesis of compounds useful for the aforementioned purposes and objectives. In a preferred embodiment, taxol is treated with mesyl chloride to prepare a taxol derivative with a contracted A-ring, which has comparable activity to taxol in a tubulin depolymerization assay, and which shows cytotoxicity against KB cells in a cell culture assay. In an alternate preferred embodiment, taxol is treated with triethyloxonium tetrafluoroborate (Meerwein's reagent) to produce a taxol derivative with an opened oxetane ring. In another alternate preferred embodiment, reaction of taxol with acetyl chloride yields a taxol derivative in which the oxetane ring is opened and the A-ring is contracted. All of the aforementioned products show in vivo activity in KB cell culture assays. Further, the preferred compounds have different in vivo activities, which makes them ideal to form a range of standards for biological testing of other compounds.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a generalized representation of a taxol compound structure.

Figure 2 is a generalized representation of a taxol compound with an opened oxetane ring.

Figure 3 is a generalized representation of a taxol compound having a contracted A-ring and an opened oxetane ring.

Figure 4 is a generalized representation of a taxol compound with a contracted A-ring.

Figure 5 is a representation of structure of an acetonide of taxol.

Figure 6 is an alternate generalized representation of a taxol compound.

Figure 7 is a generalized representation of the hydrogenated form of the compound illustrated in Figure

3.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

With reference to Figure 1, the structure of taxol (1) is represented when R1 is C(O)CH₃, R2 is OH, R3 is H, and R4 is H. The derivatives of taxol discussed herein are represented in Figures 1-7, with specific structure variables, R1-R7, defined in Table I below. For the sake of convenience, structures will be referred to by the structure numbers provided in Table I.

TABLE I

Variable Definitions For Structures Discussed Herein And Illustrated In Figures 1-7		
Structure Number	Figure Number	Variable Definitions
(1)	1	R1 = C(O)CH ₃ , R2 = OH, R3 = H, R4 = H
(2)	1	R1 = C(O)CH ₃ , R2 = H, R3 = OH, R4 = H
(3)	1	R1 = H, R2 = H, R3 = OH, R4 = H
(4)	2	R1 = C(O)CH ₃ , R2 = OH, R3 = H, R4 = H, R5 = H, R6 = H, R7 = C(O)CH ₃
(5)	2	R1 = C(O)CH ₃ , R2 = OC(O)CH ₃ , R3 = H, R4 = C(O)CH ₃ , R5 = H, R6 = H, R7 = C(O)CH ₃
(6)	2	R1 = C(O)CH ₃ , R2 = OC(O)CH ₃ , R3 = H, R4 = C(O)CH ₃ , R5 = OC(O)CH ₃ , R6 = H, R7 = C(O)CH ₃
(7)	3	R1 = C(O)CH ₃ , R2 = OC(O)CH ₃ , R3 = H, R4 = C(O)CH ₃ , R5 = OC(O)CH ₃ , R6 = OH, R7 = OC(O)CH ₃
(8)	7	R1 = C(O)CH ₃ , R2 = OC(O)CH ₃ , R3 = H, R4 = C(O)CH ₃ , R5 = OC(O)CH ₃ , R6 = OH, R7 = OC(O)CH ₃
(9)	4	R1 = C(O)CH ₃ , R2 = OSi(CH ₂ CH ₃) ₃ , R3 = Si(CH ₂ CH ₃) ₃
(10)	4	R1 = C(O)CH ₃ , R2 = OH, R3 = H
(11)	6	R1 = C(O)CH ₃ , R2 = OSi(CH ₂ CH ₃) ₃ , R3 = OH, R4 = Si(CH ₂ CH ₃) ₃
(12)	5	R1 = C(O)CH ₃ , R2 = C(O)CH ₃
(13)	6	R1 = C(O)CH ₃ , R2 = OSi(CH ₂ CH ₃) ₃ , R3 = OSO ₂ CH ₃ , R4 = Si(CH ₂ CH ₃) ₃

Due to the complexity of the taxol structure, and the incomplete understanding of the relationship between the taxol structure and its activity, it was difficult or impossible to predict how derivatives of taxol would behave; further, the complexity of the taxol structure made prediction of reaction product structures equally difficult. The present invention overcomes these problems through the synthesis of taxol derivatives with a contracted A-ring (Figure 4), with a contracted A-ring and an opened oxetane ring (Figure 3), or with an opened oxetane ring (Figure 2), which retain *in vivo* activity, although at different levels than taxol.

The taxol derivative having a contracted A-ring structure, otherwise known in a preferred embodiment as A-Nortaxol (10), is easier to perform total synthesis of than taxol since the bridged bi-cyclic portion of the diterpenoid structure has been fused. Further, it has been discovered that A-Nortaxol has an *in vitro* activity very close to that of taxol in a tubulin depolymerization assay. Thus, A-Nortaxol (10) can be used in a similar fashion to taxol (1).

It is believed that the oxetane ring (the D-ring in taxol (1)) is susceptible to ring-opening by reaction with electrophilic reagents, so initial experiments were performed using zinc bromide as an electrophile to attempt to open the oxetane ring on taxol. Taxol (1) was combined with zinc bromide in methanol at ambient temperature, but the oxetane ring was not opened, and the taxol simply underwent epimerization of the C-7 hydroxyl group and cleavage of the 10-acetyl group to yield 7-epitaxol (2) and 10-deacetyl-7-

epitaxol (3).

The relatively mild conditions created by zinc bromide in methanol were selected because it was believed that the more vigorous conditions created by other electrophiles and other solvents may result in taxol derivatives without any substantial biological activity due to destruction of part or all of the bioactive portions of the molecule. However, it was surprisingly discovered that treatment of taxol with different electrophilic groups resulted in rearrangement of different portions of the taxol structure, and that some of the various products retained *in vivo* activity. More specifically, reaction of taxol with the strong electrophile triethyloxonium tetrafluoroborate (Meerwein's reagent) resulted in a taxol derivative (4) with an opened oxetane ring. Reaction of taxol with acetyl chloride yielded a product (7) in which the oxetane ring was opened, but the A-ring was contracted. Treatment of the protected taxol (11) with mesyl chloride, followed by deprotection, resulted in a taxol derivative with a contracted A-ring alone, A-Nortaxol (10).

It was also surprisingly discovered that these rearranged taxol derivatives (4), (7), and (10) all retain varying degrees of *in vivo* activity, as determined by KB cell culture assays. A-Nortaxol (10) also showed very similar activity to taxol in a tubulin depolymerization assay, and showed higher *in vivo* activity than the acetyl chloride product (7) and the Meerwein product (4). Thus, the range of activity shown by the taxol derivatives of the present invention make them ideal for use as standards in biological testing. Further, A-nortaxol (10) and the acetyl chloride product (7) are easier to perform synthesis on than taxol since the A and B rings are fused in the products rather than bridged (note that the bridge bond between A and B rings in taxol is extremely difficult to synthesize). Thus, the present invention includes a method, which is highly useful to researchers in determining the activity of compounds in comparison to taxol (1) and taxol derivatives; the method has an additional advantage, since it is not necessary to use the limited and expensive supply of taxol as a control or standard.

In a preferred process, new compounds or production line pharmaceutical compounds, which must be tested to ensure bioactivity in comparison to taxol, are subjected to bioactivity evaluation in parallel with standards utilizing compounds (4), (7), and (10). Since the relative bioactivities of products (4), (7), and (10) are known (in comparison to one another as well as to taxol), the relative bioactivity of the compound being tested to taxol can be determined in a reliable fashion. Thus, in addition to the *in vivo* activity of compounds (4), (7), and (10), which makes the compounds useful as cytotoxic agents, they are also useful in extending the limited supply of taxol. This is because the method of the present invention replaces taxol with new compounds in testing other compounds which are used as cytotoxic agents in general, or, which are used as anti-cancer or anti-leukemic drugs in humans.

METHODS AND MATERIALS

Specific reaction methods are described in more detail in the following non-limiting examples. The methods used herein are generally described in the *Journal Of Organic Chemistry*, 51, pp. 797-802 (1986). Low resolution mass spectrometry data were obtained on a VG 7070 E-HF mass spectrometer. Exact mass measurements were performed at the Midwest Center for Mass Spectrometry, NSF Regional Instrumentation Facility (Grant Che-8211164). The term "standard work-up" in the following non-limiting examples includes extraction with a suitable solvent (usually ethyl acetate or methylene chloride), washing the extract with water, drying over magnesium sulfate or sodium sulfate, and evaporation *in vacuo*. All technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art. Other methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention.

EXAMPLE 1

Taxol (1) was reacted with zinc bromide, $ZnBr_2$ as follows: 100 mg taxol (0.117 mmol) was combined with 3.3 g $ZnBr_2$ (0.41 mmol) in 10 ml $CHCl_3:MeOH(1:4)$ and stirred for 24 hours at 40° C. Water was added to the mixture, and the standard work-up (as defined above in "Methods") yielded a white solid. The solid was purified by chromatography to obtain two compounds, which were identified as 7-epi-taxol (2) and 10-deacetyl-7-epitaxol (3) by comparison to published data (see *Journal of Natural Products*, 49, pp. 665-669 (1986), and *Journal of Natural Products*, 44, pp. 312-319 (1981); compounds (2) and (3) have been previously described by McLaughlin et al. in *Journal of Natural Products*, 44, pp. 312-319 (1981). Thus, it was necessary to try more vigorous reaction conditions to open the oxetane ring, despite the risk that the more vigorous conditions would cause other portions of the taxol structure to be altered to an extent that the resulting compound would lack any substantial bioactivity.

EXAMPLE 2

Taxol (1) was reacted with Meerwein's reagent as follows: A solution of taxol in dichloromethane was prepared by adding 100 ml taxol (0.117 mmol) to dry dichloromethane. The taxol solution in dichloromethane was cooled and stirred while 200 μ l triethyloxonium tetrafluoroborate (1 M in CH_2Cl_2) was added drop-wise from a freshly opened bottle. These conditions were maintained for 30 minutes, at which time the reaction was quenched with 3 ml of ethereal HCl (1:2 mixture of 1N HCl:ether) followed by stirring for 10 minutes. Standard work-up gave a crude solid, which was further purified by flash chromatography and PTLC to yield 53 mg (51%) of a taxol compound with an opened oxetane ring (4) (The "Meerwein product"). NMR, MS, and IR (KBr) were performed on a sample of the Meerwein product (4) and its melting point was determined, with the characterization data and NMR data presented in Tables II and III below:

TABLE II

Characterization Data For Product (4) (Resulting From Reaction Of Taxol With Meerwein's Reagent)	
Melting Point	160-164 °C (amorphous solid)
IR (KBr) cm^{-1}	1745(s), 1670(m), 1535(w), 1505(w), 1474(w), 1395(m), 1120(m), 1080(m), 1060(m)
MS(FAB) m/z	872 (MH^+ , 100), 854 (MH^+ , H_2O)
High-resolution mass spectrum	calculated (MH^+) 872.3493; observed 872.3463

TABLE III

NMR Data For Reaction Product Of Taxol With Meerwein's Reagent, $\text{C}_{47}\text{H}_{54}\text{NO}_{15}$ (4)	
Carbon Position	^1H Shift (PPM from TMS) Coupling Hertz
2	5.56 (d,6)
3	4.03 (3,6)
5	3.70 (br s)
6	a
7	4.49 (dd 4,11)
10	6.57 (s)
13	6.01 (br dd, 4,11)
14	2.45 (dd, 11,16)
	3.08 (dd, 4,16)
15	
16	1.12 (s)
17	1.12 (s)
18	2.10 (s) ^b
19	1.22 (s)
20	3.85 (ABq, 11 $\Delta\nu_{AB}$ = 86)
2'	4.70 (br s)
3'	5.92 (dd,2,9)
NH	7.19 (br d,9)
OAc	1.65 (s) 2.25 (s)
2-OBz	8.03 (m) 7.3-7.6 (m) ^c
3'-NBz	8.03 (m) 7.3-7.6 (m) ^c
3'-Ph	7.3-7.6 (m) ^c
OH	3.91 (s)

a peak concealed under signals from methyl group

b determined by decoupling experiment

c overlapping peaks

EXAMPLE 3

The Meerwein product (4) was acetylated as follows: 5 mg of the Meerwein product (4) (0.0057 mmol) was dissolved in 100 μ l of pyridine, and 300 μ l of acetyl chloride (excess) was added to the solution; the solution was kept at room temperature for 30 minutes, and warmed to 60° C for one hour, followed by quenching with water. Standard work-up yielded crude material which was purified on PTLC to yield 3 mg (55%) of chromatographically homogeneous 2',7-diacetyl-D-seco-taxol (5). Structure was confirmed with ¹H NMR.

EXAMPLE 4

2',7-diacetyl-D-seco-taxol (5) was acetylated as follows: 8 mg 2',7-diacetyl-D-seco-taxol (5) (0.008 mmol) was dissolved in 750 μ l tetrahydrofuran (THF) and to this solution dicyclohexylcarbodiimide (5 mg, 2.5 eq) 4 μ l acidic anhydride (5 eq) and a catalytic amount of pyrrolidino pyridine were added. The stirred solution was heated to 60° C for 7.5 hours, the solvent was then evaporated and the residue extracted into ethyl acetate. Standard work-up yielded a crude mixture which was purified by PTLC with 1% MeOH/CHCl₃ to yield 3 mg (38% yield at 63% conversion) of 2',5,7-triacetyl-D-seco-taxol (6). Characterization data is presented in Table IV and NMR data is presented in Table V below.

TABLE IV

Characterization Data For 2', 5, 7-Triacetyl-D-Seco-Taxol (6)	
IR cm ⁻¹	1740(s), 1720(s), 1676(m), 1625(s), 1225(s)
MS(FAB)m/z (relative intensity)	998(MH ⁺ , 13) 980(MH ⁺ , 12) 936(6) 848(23), 650(100).

TABLE V

NMR Data For 2', 5, 7-Triacetyl-D-Seco-Taxol (6)

Carbon Position	¹ H Shift (PPM from TMS) Coupling Hertz
2	5.63 (d, 5)
3	4.04 (d, 5)
5	5.26 (m) ^c
6	a
7	5.43 (br d, 9)
10	6.42 (s)
13	6.03 (m)
14	2.40 (dd, 9, 15)
	3.07 (dd, 5, 15)
15	
16	1.13 (s)
17	1.11 (s)
18	2.24 (s)
19	1.38 (s)
20	4.01 (ABq, 12 Δv _{AB} =57)
2'	5.26 (m) ^c
3'	6.12 (dd, 3, 10)
NH	7.20 (br d, 9)
OAc	1.98 (s), 2.13 (s), 2.16 (s), 2.19 (s), 2.20 (s)
2-OBz	8.20 (m)
3'-NBz	7.8 (m)
3'-Ph OH	7.2-7.6 (m) ^c

a peak concealed under signals from methyl group

b determined by decoupling experiment

c overlapping peaks

EXAMPLE 5

The Meerwein product (4) was then treated with 2,2-dimethoxypropane as follows: 6 mg of the Meerwein product (4) (0.007 mmol) and 200 μl 2,2-dimethoxypropane was combined in 500 μl dry dichloromethane along with a catalytic amount of p-toluenesulfonic acid, and stirred for one hour. Standard work-up yielded a crude product, which was further purified by PTLC to obtain 6 mg (95%) of pure acetone (12) (Figure 5). The structure of the acetone (12) was confirmed by ¹H NMR and by mass spectrometry; MS(FAB) m/z (relative intensity): 916 (MNa⁺, 100), 855 (MNa⁺-HOAc-H, 25), 832 (MNa⁺-CH₃CO-C₃H₅, 50), 761(40). ¹H NMR data is presented in Table VI below.

TABLE VI

NMR Data For Acetonide (12) Of The Meerwein Product (4)		
Position	¹ H Shift (PPM from TMS)	Coupling Hertz
2	5.34	(d, 8)
3	3.06	(d, 8)
5	4.34	(m)
6	6 _{ax} 1.95	(ddd 5, 11, 15) ^b
	6 _{eq} 2.36	(ddd 2, 5, 15)
7	4.48	(dd, 5, 11)
10	6.38	(d, 2)
13	5.68	(d, 2)
14	2.57	(AB part of ABX, 9, 14 Δv _{AB} =62)
15		
16	4.67	(s), 4.75 (s)
17	1.63	(s)
18	1.67	(s) ^b
19	1.62	(s)
20	4.15	(ABq, 12, Δv _{AB} =26)
2'	4.50	(d, 3)
3'	5.60	(dd, 3, 8)
NH	6.98	(br d, 8)
OAc	1.83	(s), 2.17 (s)
2-OBz	8.10	(m)
	7.3-7.4	(m) ^c
3'-NBz	7.73	(m)
	7.3-7.4	(m) ^c
3'-Ph	7.3-7.4	(m) ^c
OH		
Other	1.30	(s) ^d
	1.33	(s)

a peak concealed under signals from methyl group

b determined by decoupling experiment

c overlapping peaks

d signals of the methyl groups of the acetonide

EXAMPLE 6

Taxol (1) was reacted with acetyl chloride under the following conditions: 200 mg taxol (0.23 mmol) was dissolved in 2 ml acetyl chloride, and the solution was refluxed for one hour. The reaction was quenched with ice water and ethyl acetate, and stirred for 30 minutes. Standard work-up yielded a white solid. Recrystallization of the white solid from ethyl acetate and hexanes yielded acetylated A-Nor-D-seco-taxol (7) (See Figure 3) as white needles (156 mg, 68%). Characterization data for acetylated A-Nor-D-seco-taxol (7) is provide in Table VII and NMR data is provided in Table VIII below.

TABLE VII

Characterization Data For Acetylated A-Nor-D-seco-taxol (7)	
Melting Point	140-142 °C
IR(CHCl ₃) cm ⁻¹	1750(s), 1660(m), 1606(m), 1372(m), 1282(m), 1156(m)
MS(FAB)m/z	1002(MNa ⁺ , 35), 676(MNa ⁺ -side-chain, 15), 616 (675-HOAc, 30), 554(676-PhCOOH, 20), 494(616-PhCOOH, 30), 411(24), 3722(4), 177(100)
High resolution mass spectrum	calculated for C ₅₃ H ₅₇ NO ₁₇ Na(MNa ⁺) = 1002.3524; observed 1002.3557

TABLE VIII

NMR Data For Acetylated A-Nor-D-seco-taxol (7)		
Carbon Position	¹ H Shift (PPM from TMS)	Coupling Hertz
2	5.35 (d, 7)	

	3	3.54 (d, 7)
	5	5.28 (br s)
	6	a
5	7	5.54 (dd, 4, 13)
	10	6.38 (s)
	13	5.67 (t, 7)
	14	2.64 (m)
	15	
10	16	4.69 (s), 4.82 (s)
	17	1.62 (s)
	18	1.82 (s)
	19	1.53 (s)
	20	4.15 (ABq, 12
15		$\Delta\nu_{AB}=55$)
	2'	5.52 (d, 2)
	3'	6.02 (dd, 2, 9)
	NH	7.03 (d, 9)
	OAc	1.85 (s), 2.00 (s),
20		2.13 (s), 2.17 (s),
		2.22 (s)
	2-OBz	7.93 (m),
		7.2-7.9 (m) ^c
	3'-NBz	7.85 (m)
25		7.2-7.9 (m) ^c
	3'-Ph	7.2-7.9 (m) ^c
	OH	3.75 (s)
	Other	

- a peak concealed under signals from methyl group
 b determined by decoupling experiment
 c overlapping peaks

35 EXAMPLE 7

The Meerwein product (4) was treated with acetyl chloride as follows: 50 mg of the Meerwein product (4) (0.057 mmol) was dissolved in CHCl_3 ; 3 mg pyrrolidinopyridine (an amount sufficient to catalyze the reaction) and excess acetyl chloride (5eq) were added to the solution. Excess triethylamine was then added dropwise to the stirred solution at room temperature. A red color developed, which disappeared when more acetyl chloride was added; additional triethylamine and acetyl chloride were added until a single major product was obtained. The reaction was stopped after a total reaction time of 5 hours by addition of 3 ml of water and stirring for 30 minutes. Standard work-up was followed with an additional wash with 3N HCl. The mixture of products was subjected to PTLC to obtain the major component as 10 mg of a white solid (18%).
¹H NMR, mass spectrometry, and infrared spectrometry confirmed that this compound was identical to the acetyl product of Example 6, acetylated A-Nor-D-seco-taxol (7)

EXAMPLE 8

The acetylated A-Nor-D-seco-taxol (7) was hydrogenated to its dihydro derivative (8) (Figure 7) as follows: 24 mg of the acetyl chloride product (7) (0.023 mmol) was dissolved in 2.5 ml ethyl acetate, and hydrogenated over Pd/H_2 . After 24 hours, the catalyst was filtered off, and the solvent was evaporated to yield a crude solid, which consisted of product and unreacted starting material; the unreacted starting material was not separable from the product. The crude product was dissolved in methylene chloride and treated with 5 mg m-chloroperbenzoic acid (58%) at room temperature for 3 hours; this converted the starting material to its separable epoxide. The solvent was evaporated and the residue subjected to PTLC with 4% $\text{MeOH}/\text{CHCl}_3$ to yield 8 mg pure hydrogenated product (8) (35%) along with 11 mg of a mixture of diastereomeric epoxides. The hydrogenated product (8) was recrystallized from ethyl acetate and hexanes.

Characterization data is presented in Table I below and NMR data is presented in Table IX below.

TABLE IX

Characterization Data for Hydrogenated Acetylated A-Nor-D-Seco-Taxol (8)	
Melting Point	148-150° C
IR (KBr) cm^{-1}	1740(s), 1720(m), 1640(m), 1220(m), 910(m)
MS(FAB) m/z (relative intensity)	1004(MNa^+ , 100), 962($\text{MNa}^+ - \text{C}_3\text{H}_6$, 10), 944($\text{MNa}^+ - \text{HOAc}$, 15)

TABLE X

NMR Data For Hydrogenated Acetylated A-Nor-D-Seco-Taxol (8)	
Carbon Position	^1H Shift (PPM from TMS) Coupling Hertz
2	5.17 (d, 8)
3	3.68 (d, 8)
5	4.94 (br d, 11)
6	a
7	5.54 (dd, 5, 11)
10	6.36 (s)
13	5.72 (m)
14	2.48 (AB part of ABX, 14, 16 $\Delta\nu_{AB}=65$)
15	1.60 (m)
16	0.76 (d, 7)
17	0.78 (d, 7)
18	1.83 (s)
19	1.53 (s)
20	4.11 (ABq, 11 $\Delta\nu_{AB}=78$)
2'	5.40 (d, 3)
3'	5.95 (dd, 3, 8)
NH	7.02 (d, 8)
OAc	1.17 (s), 1.18 (s), 2.00 (s), 2.13 (s), 2.17 (s)
2-OBz	7.91 (m), 7.2-7.6 (m) ^c
3'-NBz	7.83 (m) 7.2-7.6 (m) ^c
3'-Ph	7.2-7.6 (m) ^c
OH	
Other	

a peak concealed under signals from methyl group

b determined by decoupling experiment

c overlapping peaks

EXAMPLE 9

Taxol was reacted with imidazole and triethylsilylchloride as follows: 238 mg of solid imidazole (10 eq) was added to a solution of 200 mg taxol (0.234 mmol) in 2.5 ml DMF; 196 μ l triethylsilylchloride (10 eq) was then added to the stirred solution while the solution was at room temperature, followed by warming the solution to 45-50 °C. After 2 hours, the solution was diluted with water, and extracted with ethyl acetate. The crude solid obtained after evaporation of the solvent was purified on a silica gel flash column to yield 242 mg (96%) of pure 2',7-bis (triethylsilyl)taxol (11) (Figure 6). Characterization data is presented in Table XI and NMR data is presented in the Table XII below.

TABLE XI

Characterization Data For 2',7-bis(triethylsilyl)taxol (11)	
Melting Point	122-123°C
IR cm^{-1}	1740(s), 1720(s), 1660(s), 1640(m), 1240(s), 810(m)
MS(FAB) m/z (relative intensity)	1104(MNa^+ , 100), 1003(30), 981(MNa^+ -PhCOOH, 10)

TABLE XII

NMR Data For 2',7-bis(triethylsilyl)taxol (11)	
Carbon Position	^1H Shift (PPM from TMS) Coupling Hertz
2	5.71 (m)
3	3.88 (d,6)
5	4.94 (br d, 11)
6	a
7	4.48 (dd,6,14)
10	6.44 (s)
13	6.20 (m)
14	a
15	
16	1.22 (s)
17	1.18 (s)
18	2.02 (s)
19	1.70 (s)
20	4.27 (ABq, 11 $\Delta\nu_{AB} = 63$)
2'	4.71 (d, 3)
3'	5.71 (m)
NH	7.12 (d,9)
OAc	2.17 (s), 2.55 (s)
2-OBz	8.12 (m) 7.2-7.5 (m) ^c
3'-NBz	7.79 (m) 7.2-7.5 (m) ^c
3'-Ph	7.2-7.5 (m) ^c
OH	
Other	0.48 (6H,m) ^c 0.57 (6H,m) 0.81 (9H,t,8) 0.91 (9H,t,8)

a peak concealed under signals from methyl group

b determined by decoupling experiment

c overlapping peaks

EXAMPLE 10

2',7-bis(triethylsilyl)taxol (11) was treated with triethylamine and mesyl chloride (methane sulfonyl chloride) as follows: 30 mg of 2',7-bis(triethylsilyl)taxol (11) (0.028 mmol) was added to 3 ml of dry methylene chloride, and cooled to -15 °C under argon. 600 μ l of triethylamine (154 eq) was added, followed by addition of 300 μ l of mesyl chloride (138 eq) in 1 ml methylene chloride, with the additions made over a

5 minute period. The mixture was allowed to warm to -5 to 0° C, and this temperature was maintained for a total reaction time of 2.5 hours, at which time 50% conversion of the starting material to 2',7-bis-(triethylsilyl)-A-Nortaxol (9) (Figure 4) was observed. The solution was cooled again to -15° C, and 1 ml triethylamine and 500µl mesyl chloride were added; this latter procedure was repeated one additional time.

The reaction was then stopped by adding 2 ml triethylamine, 5 ml water, and 5 ml ethyl acetate. Standard work-up yielded crude material, which was purified by PTLC to give 6 mg (20%) of 2',7-bis(triethylsilyl)-A-Nor-taxol (9) (Figure 4) along with 2 mg starting material and 2 mg 7-triethylsilyl-A-Nor-taxol. It is believed that the mesylate (13) (Figure 6) is formed as an intermediate between (11) and (9). Mass spectrometry data, m/z, are as follows: 1086(MNa⁺, 45), 1064(MH⁺, 75), 1005(MH⁺-OAc, 25), 975(MH⁺-OAc-CH₂O, 15), 963(MH⁺-OAc-C₃H₆, 15), 820(MH⁺-PhCOOH, PhCONH₂-H, 100). NMR data is presented in Table XIII below:

TABLE XIII

NMR Data For 2',7-bis(triethylsilyl)-A-Nor-taxol(9)	
Carbon Position	¹ H Shift (PPM from TMS) Coupling Hertz
2	5.54 (d,8)
3	3.53 (d,8)
5	5.02 (d,8)
6	1.90 (m)
7	4.53 (dd,8,5)
10	6.39 (s)
13	5.81 (br t,7)
14	2.40 (m) 2.60 (m)
15	
16	4.66 (s), 4.75 (s)
17	1.65 (s)
18	1.62 (s)
19	1.73 (s)
20	4.15 (ABq, 12) $\Delta \nu_{AB} = 26$
2'	4.60 (d,2)
3'	5.64 (dd,2,11)
NH	7.15 (d,11)
OAc	2.02 (s), 2.40 (s)
2-OBz	8.01 (m) 7.2-7.5 (m) ^c
3'-NBz	7.70 (m) 7.2-7.5 (m) ^c
3'-Ph	7.2-7.5 (m) ^c
OH	
Other	0.38 (6H,m) ^a 0.60 (6H,m) 0.75 (9H,t, 8) 0.90 (9H,t, 8)

a peak concealed under signals from methyl group

b determined by decoupling experiment

c overlapping peaks

e signals of the ethyl groups of the TES groups

EXAMPLE 11

2',7-bis(triethylsilyl)-A-Nor-taxol (9) was treated with pyridinium hydrofluoride as follows: 67 ml 2',7-bis-(triethylsilyl)-A-Nor-taxol (9) (0.07 mmol) was dissolved in 1 ml dry THF under argon. The solution was cooled to 0° C and 100µl pyridinium hydrofluoride (70% in pyridine) was added. After 3 hours the cooling bath was removed, and the reaction was allowed to proceed for an additional 45 hours at room temperature. The reaction was quenched with 2 ml aqueous pyridine (10% v/v pyridine). Standard work-up resulted in a crude solid which was purified by PTLC with 8% MeOH/CHCl₃, as the eluent. A-nor-taxol (10) was obtained as a white solid in 55% yield (29 mg). (Note that deprotection of the silyl ether (9) yielded a mixture of products when tetrabutylammonium fluoride was used instead of pyridinium hydrofluoride.) Mass spectrometry data, m/z (relative intensity), as follows: 836(MH⁺, 100), 776(MH⁺-HOAc, 30), 551(836-side-chain-H, 10), 307(20); High resolution mass spectrum calculated for C₄₇H₅₀NO₁₃(MH⁺) was 836.3282, observed

836.3272. ¹H NMR is presented in Table XIV below.

TABLE XIV

¹ H NMR Data For A-Nor-Taxol (10)	
Carbon Position	¹ H Shift (PPM from TMS) Coupling Hertz
2	5.49 (d,8)
3	3.48 (d,8)
5	5.04 (d,8)
6	1.86 (dd,11,15); 2.59 (ddd,8,9,15)
7	4.63 (dd,9,11)
10	6.32 (s)
13	5.71 (m) ^c
14	2.04 (dd,8,13) 2.42 (dd,13,6)
15	
16	4.69 (br s) 4.76 (br s)
17	1.59 (s)
18	1.61 (s)
19	1.64 (s)
20	4.24 (ABq, 8, Δν _{AB} = 34)
2'	4.67 (d,2)
3'	5.71 (dd,11,2) ^c
NH	6.89 (d,11)
OAc	2.17 (s), 2.36 (s)
2-OBz	8.10 (m) 7.2-7.6 (m) ^c
3'-NBz	7.66 (m) 7.2-7.6 (m) ^c
3'-Ph	7.2-7.6 (m) ^c
OH	
Other	

a peak concealed under signals from methyl group

b determined by decoupling experiment

c overlapping peaks

BIOLOGICAL TESTING OF TAXOL DERIVATIVES

Taxol (1), the Meerwein product (4), the acetyl chloride product (7), and A-nor-taxol (10) were tested in KB cell culture assays. All of the samples showed *in vivo* activity as shown in Table XV below (note that an ED₅₀ (μg/ml) of less than 4.0 indicates *in vivo* activity). Taxol has been shown to have great potential as an anti-cancer drug, so that these compounds can be used in a similar fashion to taxol as cytotoxic agents.

TABLE XV

Bioactivity Of Modified Taxols In KB Cell Culture	
Compound	ED ₅₀ (μg/ml) cell culture
Taxol (1)	0.00001
Meerwein product (4)	2.3
Acetyl Chloride product (7)	2.5
A-Nortaxol (10)	2.0

Taxol (1), the Meerwein product (4), the acetyl chloride product (7), and A-Nor-Taxol (10) were also subjected to tubulin depolymerization assays with the results presented in Table XVI below.

TABLE XVI

In Vitro Activity In Tubulin Depolymerization Assay	
Compound	ID ₅₀ (μ M) in tubulin depolymerization assay
Taxol (1)	0.3
Meerwein product (4)	>6.3
A-Nortaxol (10)	0.9

Note that A-Nor-Taxol (10) has an activity which is very similar to taxol in the tubulin depolymerization assay. Since the A-Nor-Taxol (10) structure has a fused A-ring, A-Nor-Taxol (10) is a more likely candidate for total synthesis than taxol. Further, because A-Nor-taxol has been found to be cytotoxic, and to have an in vitro activity close to taxol's in a tubulin depolymerization assay, it is believed that A-Nor-taxol (10) may have bioactivity mechanisms very similar to taxol's (1).

With reference to Table XV, note that the in vivo activity of A-Nor-Taxol (10), the Meerwein product (4), and the acetyl chloride product (7) are 2.0, 2.3, and 2.5 respectively. This range of in vivo activities makes these compounds useful as standards for testing other compounds, eliminating the need for taxol standards. Since, the relative in vivo activity of these compounds is now known, the bioactivity of an unknown compound can be determined with respect to these compounds without having to refer to any published data or without use of taxol as a standard.

A preferred process for testing the bioactivity of unknown compounds, or for testing taxol-derived pharmaceuticals, includes treatment of an unknown sample in a bioactivity test, and simultaneous treatment under the same conditions of at least one, preferably of all three, of the following products: A-Nor-Taxol (10), the Meerwein product (4), and the acetyl chloride product (7). By comparison of results for the unknown sample with the results for the products (4), (7), and (10), the in vivo activity of the unknown can be determined.

Due to the bioactivity demonstrated for some of the compounds of the present invention, it is believed that compounds (4)-(12) will be useful directly as, or as intermediates for, antineoplastic, antileukemic and anticancer prodrugs or drugs, and their use as such is considered to be a part of the present invention.

Contemplated equivalents of the rearranged taxol compounds of the present invention include rearranged taxol compounds having a contracted A-ring, an opened oxetane ring, or a contracted A-ring and an opened oxetane ring, which have one or more side chain or ring substituents substituted with a non-interfering group (e.g., a substituted group which does not seriously alter the desirable properties of the rearranged taxol compounds of the present invention), such as, but not limited to, substitution of -H, -OH, -OR, -NR, -Ar, -OC(O)CH₃, Ph-O-, Ph-N-, or =O for another non-interfering moiety.

From the above teachings, it is apparent that many modifications and variations of the present invention are possible. It is therefore to be understood that the invention may be practiced otherwise than as specifically described.


Claims

1. A rearranged taxol compound having the following structure:

5

10

15



20


wherein R1 is C(O)CH_3 , R2 is H or $\text{Si}(\text{CH}_2\text{CH}_3)_3$, and R3 is H or $\text{Si}(\text{CH}_2\text{CH}_3)_3$.

2. A taxol compound having the following structure:

25

30

35



40


wherein R1 is C(O)CH_3 , R2 is H or $\text{OSi}(\text{CH}_2\text{CH}_3)_3$, R3 is OSO_2CH_3 , and R4 is H or $\text{Si}(\text{CH}_2\text{CH}_3)_3$.

3. A rearranged taxol compound having the following structure:

45

50

55



wherein R1 is H or C(O)CH₃, R2 is H or C(O)CH₃.

4. A process for producing taxol compounds with an A-ring contracted to five members, comprising the steps of:
 - (a) protecting hydroxylated positions on taxol; and
 - (b) reacting said protected taxol with a mesyl halide.
5. A process according to claim 4, further comprising the step of deprotecting said protected taxol to form A-Nor-taxol.
6. A process for producing taxol compounds with an A-ring contracted to five members, comprising the steps of:
 - (a) reacting taxol with a trialkylsilyl halide to form a taxol silyl ether; and
 - (b) reacting taxol silyl ether with a mesyl halide.
7. A process according to claim 6, further comprising the step of: (c) following step (b) removing silyl groups on said taxol silyl ether to yield A-Nor-taxol.
8. The process of claim 7, wherein step (c) is carried out by reacting said silylated A-Nor-taxol with pyridinium hydrofluoride.
9. The process of claim 6, 7 or 8, wherein said steps (a) and (b) are carried out in basic conditions.
10. The process of any one of claims 6, to 9, wherein said trialkylsilyl halide is triethylsilyl chloride, and said mesyl halide is mesyl chloride.
11. A process for producing rearranged taxol compounds comprising the reaction of taxol with an oxonium compound to form a taxol compound with an opened oxetane ring; and reacting said taxol compound with an opened oxetane ring with dimethoxypropane in the presence or p-toluenesulfonic acid.
12. The process of claim 11, wherein said oxonium compound is triethyloxonium tetrafluoroborate.
13. A process for determining the bioactivity of a compound, comprising treatment of an unknown compound in a bioactivity test and treatment under the same conditions of at least one of the compounds selected from the group comprised of A-Nor-taxol, a taxol compound with an opened oxetane ring, and a taxol compound with an opened oxetane ring and a five member A-ring fused to the B-ring.
14. A cytotoxic agent comprising a taxol compound having an A-ring contracted to five members.
15. The cytotoxic agent of claim 14, wherein said taxol compound has an intact oxetane D-ring.




FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number

EP 92 30 1652

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
A	JOURNAL OF THE AMERICAN CHEMICAL SOCIETY, vol. 93, no. 9, 5 May 1971, Washington, DC, US, pages 2325 - 2327; M.C. WANI ET AL.: 'Plant antitumour agents. VI. The isolation and structure of taxol, a novel antileukaemic and antitumour agent from Taxus brevifolia' * page 2325 *	1, 14	C07D305/14 C07D317/70 A61K31/335
P, A	JOURNAL OF ORGANIC CHEMISTRY, vol. 56, no. 17, 16 August 1991, Washington, DC, US, pages 5114 - 5119; G. SAMARANAYAKE ET AL.: 'Modified taxols. 5. Reaction of taxol with electrophilic reagents and preparation of a rearranged taxol derivative with tubulin assembly activity' * see the whole document, especially compounds 8, 20, 21, 22; page 5117, table I *	1-12, 14, 15	TECHNICAL FIELDS SEARCHED (Int. Cl.5) C07D
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 26 MAY 1992	Examiner RUSSELL F. ENGLISH
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons * : member of the same patent family, corresponding document	